

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number: **0 596 254 A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **93115917.2**

(22) Date of filing: **01.10.93**

(51) Int. Cl.⁵: **C07C 251/52, A01N 37/50,
C07C 251/48, C07C 251/54,
C07C 251/58, C07C 255/62,
C07C 251/88, C07C 251/68,
C07D 213/61, C07D 317/54,
C07D 333/28, C07D 277/32**

(30) Priority: **02.10.92 JP 265193/92
05.11.92 JP 296140/92
15.07.93 JP 175497/93**

(43) Date of publication of application:
11.05.94 Bulletin 94/19

(84) Designated Contracting States:
BE CH DE DK ES FR GB IT LI NL

(71) Applicant: **MITSUBISHI KASEI CORPORATION
5-2, Marunouchi 2-chome
Chiyoda-ku
Tokyo(JP)**


(72) Inventor: **Oda, Masatsugu**

**42-24, Kamiseya-cho,
Seya-ku
Yokohama-shi, Kanagawa(JP)
Inventor: Katsurada, Manabu
2340-4, Shindo
Sagamihara-shi, Kanagawa(JP)
Inventor: Tomita, Hirofumi
B-501, 3-8-3, Minamitsukushino
Machida-shi, Tokyo(JP)**

(74) Representative: **TER MEER - MÜLLER -
STEINMEISTER & PARTNER
Mauerkircherstrasse 45
D-81679 München (DE)**

(54) **Methoxyiminoacetic acid derivative and agricultural/horticultural fungicide containing the same as active ingredient.**

(57) A methoxyiminoacetic acid derivative represented by the following formula (I):



(I)

wherein X represents a hydrogen atom, a halogen atom, an alkyl group having 1 to 4 carbon atoms or an alkoxy group having 1 to 4 carbon atoms; A represents a methoxy group or a methylamino group; when A is a methoxy group, B represents -O-CO- or -N=C(R¹)- and when A is a methylamino group, B represents -O-CR¹R²-, wherein R¹ and R² independently represent a hydrogen atom, an alkyl group having 1 to 4 carbon atoms or a trifluoromethyl group; and Ar represents an optionally substituted aryl group or an optionally substituted heteroaryl group, and an agricultural/horticultural fungicide containing the same as an active ingredient.

EP 0 596 254 A1


26

This invention relates to a novel methoxyiminoacetic acid derivative and an agricultural/horticultural fungicide containing the same as an active ingredient.

It has been known that certain methoxyiminoacetic acid derivatives have biological activities including fungicidal activities. For example, a compound of the formula:

5

10




15

is described in EP 398692. Further a compound of the formula:

20

25




is described in WO92/13830 and EP 463488. Furthermore, a compound of the formula:

30

35


40



is described in EP 499823. Also, a compound of the formula:

45

50




is described in EP515901.

However, these compounds are not always satisfactory as an agricultural/horticultural fungicide, as will be shown in Test Examples hereinafter.

Under these circumstances, the present inventors have paid their attention to these methoxyiminoacetic acid derivatives and conducted extensive studies thereon. As a result, it has successfully been found out that a methoxyiminoacetic acid derivative having a specific structure has a potent fungicidal activity as well as an excellent systemic and residual activity for plants, thus completing the present invention.

Accordingly, the gist of the present invention resides in a methoxyiminoacetic acid derivative represented by the following formula (I):



wherein X represents a hydrogen atom, a halogen atom, an alkyl group having 1 to 4 carbon atoms or an alkoxy group having 1 to 4 carbon atoms; A represents a methoxy group or a methylamino group; when A is a methoxy group, B represents -O-CO- or -N=C(R¹)- and when A is a methylamino group, B represents -O-CR¹R²-, wherein R¹ and R² independently represent a hydrogen atom, an alkyl group having 1 to 4 carbon atoms, a cyano group or a trifluoromethyl group; and Ar represents an optionally substituted aryl group or an optionally substituted heteroaryl group, and an agricultural/horticultural fungicide containing the same as an active ingredient.

Now, the present invention will be described in detail.

The methoxyiminoacetic acid derivative of the present invention is the one represented by the above formula (I). In the above formula (I), X represents a hydrogen atom; a halogen atom (for example, fluorine, chlorine, bromine); an alkyl group having 1 to 4 carbon atoms (for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl); or an alkoxy group having 1 to 4 carbon atoms (for example, methoxy, ethoxy, iso-propoxy, n-butoxy). It preferably represents a hydrogen atom or a halogen atom, still preferably a hydrogen atom.

A represents a methoxy group or a methylamino group. It preferably represents a methylamino group.

When A is a methoxy group, B represents -O-CO- or -N=C(R¹)-. When A is a methylamino group, B represents -O-CR¹R²-. R¹ and R² independently represent a hydrogen atom; an alkyl group having 1 to 4 carbon atoms (for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl); a cyano group; or a trifluoromethyl group. It preferably represents a hydrogen atom, a cyano group or a methyl group.


Ar represents an aryl group (for example, phenyl, naphthyl) which may be optionally substituted by the following groups; or a heteroaryl group (for example, pyridyl, thienyl, thiazolyl) which may be optionally substituted by the following groups. It preferably represents a phenyl group which may be optionally substituted by the following groups, a naphthyl group, a thienyl group which may be optionally substituted by the following groups, or a thiazolyl group which may be optionally substituted by the following groups.

Examples of the substituents for the above-mentioned aryl group include a cyano group; a halogen atom (for example, fluorine, chlorine, bromine); an alkyl group having 1 to 6 carbon atoms (for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl); an alkenyl group having 2 to 4 carbon atoms (for example, ethenyl, propenyl) optionally substituted by a halogen atom; a haloalkyl group having 1 to 4 carbon atoms (for example, trifluoromethyl, difluoromethyl, trichloromethyl, dichlorodifluoroethyl); an alkoxy group having 1 to 6 carbon atoms (for example, methoxy, ethoxy, iso-propoxy, n-butoxy) optionally substituted by a halogen atom or a cycloalkyl group having 3 to 6 carbon atoms; an alkylcarbonyloxy group having 1 to 7 carbon atoms (for example, acetoxy, propionyloxy, pivaloyloxy) optionally substituted by a halogen atom; an acylamino group having 1 to 7 carbon atoms (for example, acetoamino, propionylamino) optionally substituted by a halogen atom; an alkylthio group having 1 to 6 carbon atoms (for example, methylthio, ethylthio, iso-propylthio, n-butylthio) optionally substituted by a halogen atom; an aryl group (for example, phenyl) optionally substituted by an alkyl group having 1 to 4 carbon atoms or a halogen atom; an aryloxy group (for example, phenoxy) optionally substituted by an alkyl group having 1 to 4 carbon atoms or a halogen atom; an alkylsulfonyloxy group having 1 to 6 carbon atoms (for example, methanesulfonyloxy or ethanesulfonyloxy) optionally substituted by a halogen atom; an alkenyloxy group having 2 to 6 carbon atoms (for example, propenyloxy) optionally substituted by a halogen atom; and an alkynyloxy group having 2 to 6 carbon atoms (for example, propargyloxy). From among these substituents, those adjacent to each other may be combined together to give, for example, a methylenedioxy or ethylenedioxy group and form a fused ring together with an aryl group. The number of substituents is from 1 to 5, preferably from 1 to 2. When Ar has two or more substituents, they may be the same or different each other. Preferable examples of substituents for an aryl group include an alkyl group having 1 to 4 carbon atoms, a halogen atom, an

alkoxy group having 1 to 4 carbon atoms which may be optionally substituted by a halogen atom (preferably fluorine), an acylamino group having 1 to 4 carbon atoms which may be optionally substituted by a halogen atom (preferably fluorine), an alkylthio group having 1 to 3 carbon atoms, an alkylsulfonyloxy group having 1 to 3 carbon atoms which may be optionally substituted by a halogen atom (preferably fluorine) and a trifluoromethyl group.

Examples of the substituents for the above-mentioned heteroaryl group include a cyano group; a halogen atom (for example, fluorine, chlorine, bromine); an alkyl group having 1 to 6 carbon atoms (for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl); a haloalkyl group having 1 to 4 carbon atoms (for example, trifluoromethyl, difluoromethyl, trichloromethyl, dichlorodifluoroethyl); and an alkoxy group having 1 to 6 carbon atoms (for example, methoxy, ethoxy, iso-propoxy, n-butoxy) optionally substituted by a halogen atom or a cycloalkyl group having 3 to 6 carbon atoms. The number of substituents, which may be the same or different each other, is from 1 to 2. Among these substituents, an alkyl group having 1 to 4 carbon atoms, a halogen atom and a trifluoromethyl group may be cited as preferable ones.

The compounds of the present invention are each a novel one and can be prepared, for example, in accordance with the following reaction scheme:



[wherein X, R¹, R² and Ar are as defined in the above formula (I)].

The compounds represented by the above formulae (II) and (I-a) can be prepared by, respectively, reacting the benzaldehyde derivatives of the above formulae (III) and (V) with a hydroxylamine hydrochloride and reacting the oxime derivatives (IV) and (VI) thus obtained with the corresponding benzyl halide

derivative or benzoyl halide derivative in the presence of an appropriate base in an inert solvent (for example, diethyl ether, tetrahydrofuran, dimethylformamide, dimethylsulfoxide, methylene chloride, dichloroethane).


Examples of the base to be used in the above reaction include an alkali metal hydride (for example, sodium hydride); an alkali metal alcoholate (for example, sodium methylate); an alkali metal carbonate (for example, potassium carbonate); an alkali metal hydroxide (for example, potassium hydroxide); a tertiary amine (for example, N-methylmorpholine, triethylamine); and an aromatic base (for example, pyridine, picoline).

In some cases, the compound of the above formula (II) can be directly obtained by reacting the benzaldehyde derivative (III) with o-substituted hydroxylamine in an inert solvent such as an alcohol.

The compound of the above formula (I-b) can be obtained by reacting the benzaldehyde derivative represented by the above formula (V) with the corresponding hydrazone derivative in an inert solvent such as an alcohol.

The above-mentioned compounds of formulae (III) and (V) as starting materials can be produced in accordance with the method described in EP398692, EP499823 or the like.

In some cases, the compound of the above formula (II) can be prepared in accordance with the following reaction scheme:



[wherein R^1 , R^2 and Ar are as defined in the above formula (I) and R^3 represents an alkyl group having 1 to 10 carbon atoms].

The compound of the above formula (II) can be obtained by reacting an ester derivative (VII) with methylamine in an inert solvent such as an alcohol.

The compounds of the above formulae (II), (I-a) and (I-b) each exists as isomers at the methoxyimino moiety. Each isomer can be separated each other from the mixture of isomers which is obtained usually, by a conventional manner such as column chromatography. Each of E-, Z-mixture or Z-isomer can be converted to the E-isomer which shows high activities, by treating with an acid (for example, hydrochloric acid, sulfuric acid, methanesulfonic acid) in an alcohol solvent (for example, methanol, ethanol).

The compounds of the present invention thus obtained are each a novel one having an excellent fungicidal activity. They exert excellent preventive effects on various phytopathogenic fungi, which makes them useful as an agricultural/horticultural fungicide.

For example, these compounds exert high activity on rice blast (*Pyricularia oryzae*), rice sheath blight (*Rhizoctonia solani*), wheat powdery mildew (*Erysiphe graminis f. sp. tritici*) and barley powdery mildew (*E. graminis f. sp. hodei*), various leaf rusts of wheat and barley (e.g., *Puccinia recondita*), gray mold of vegetables and fruit trees (*Botrytis cinerea*) and late blight of various crops (*Phytophthora infestans*). Further, they have prolonged residual activity and excellent systemic action in plants, which makes them highly useful as an agricultural/horticultural fungicide.

When the compound of the present invention is to be used as an agricultural/horticultural fungicide, it may be applied as such. However, it is preferable to formulate said compound into, for example, emulsifiable concentrate, wettable powder, dust or granules by blending with adjuvants in a conventional manner to thereby ensure the effective dispersion of the active ingredient at the application.

When the agricultural/horticultural fungicide according to the present invention is to be formulated into an emulsifiable concentrate, 10 to 80 parts by weight (hereinafter referred to as "parts") (preferably 10 to 70 parts) of the compound of the present invention, 10 to 90 parts (preferably 20 to 80 parts) of a solvent and 3 to 20 parts (preferably 5 to 15 parts) of a surfactant are mixed together at an appropriate ratio. At the

usage, the obtained mixture is diluted with water to a definite concentration and applied by, for example, spraying.

When the agricultural/horticultural fungicide of the present invention is to be used as a wettable powder, 5 to 80 parts (preferably 10 to 70 parts) of the compound of the present invention, 10 to 90 parts (preferably 20 to 80 parts) of a filler and 1 to 20 parts (preferably 3 to 15 parts) of a surfactant are mixed together at an appropriate ratio. At the usage, the obtained mixture is diluted with, for example, water to a definite concentration and applied, similar to the case of the emulsifiable concentrate.

When the agricultural/horticultural fungicide of the present invention is to be used as a dust, 0.1 to 10 parts (preferably 1 to 5 parts) of the compound of the present invention is uniformly mixed with 90 to 99.9 parts (preferably 95 to 99 parts) of a filler (for example, kaolin, bentonite, talc).

The agricultural/horticultural fungicide of the present invention may further contain other active ingredients such as bactericides, insecticides and miticides, so long as the effects of the active ingredient of the present invention are not deteriorated thereby.

The agricultural/horticultural fungicide of the present invention can be suitably used either in foliar application or in submerged application. In the case of foliar application, the agricultural/horticultural fungicide is usually formulated into an emulsifiable concentrate or a wettable powder and diluted with water so as to give a concentration of the active ingredient of from 10 to 1,000 ppm. Then it is applied at a ratio of 100 to 5000 l per 1 ha.

To further illustrate the present invention in greater detail, the following Examples will be given. However, it is to be understood that the present invention is not restricted thereto but various changes may be restored within the scope thereof.

SYNTHESIS EXAMPLE 1

Synthesis of N-methyl-2-[2-{3-(trifluoromethyl)benzyloxyiminomethyl}phenyl]-2-methoxyiminoacetamide (compound No. 1 in Table 1):

To a solution of 0.53 g of methyl 2-[2-{3-(trifluoromethyl)benzyloxyiminomethyl}phenyl]-2-methoxyiminoacetate in 5 ml of methanol, was added 5 ml of a 40 % methylamine/methanol solution and the mixture was stirred at room temperature overnight. After the completion of the reaction, the solvent was distilled off and the residue was recrystallized from ethyl acetate/hexane (1 : 9). Thus 0.56 g of the title compound was obtained (quantitative yield).

The compound No. 2 in Table 1 and the compound No. 78 in Table 2 were synthesized by repeating the above-mentioned procedure except altering the starting material.

SYNTHESIS EXAMPLE 2

Synthesis of N-methyl-2-[2-{3-chlorobenzyloxyiminomethyl}phenyl]-2-methoxyiminoacetamide (compound No. 3 in Table 1):

To a solution of 1 g (4.26 mmol) of N-methyl-2-[2-(hydroxyiminomethyl)phenyl]-2-methoxyiminoacetamide and 0.62 g (4.5 mmol) of potassium carbonate in 10 ml of DMF, was added 0.69 g (4.29 mmol) of 3-chlorobenzyl chloride and the mixture was stirred under heating at 110 °C for 3 hours. After cooling, the reaction mixture was poured into water, extracted with ethyl acetate, successively washed with water and a saturated sodium chloride solution and dried over anhydrous sodium sulfate. After concentration in vacuo, the residue was chromatographed over SiO₂ to give 0.9 g of the title compound (yield: 58.8 %).

The compounds No. 4 to No. 8 and No. 11 to No. 14 in Table 1 were synthesized by repeating the above-mentioned procedure except altering the starting material.

SYNTHESIS EXAMPLE 3

Synthesis of N-methyl-2-[2-{ α -methyl-4(trifluoromethyl)benzyloxyiminomethyl}phenyl]-2-methoxyiminoacetamide (compound No. 24 in Table 1):

To a solution of 0.6 g (2.7 mmol) of N-methyl-2-(2-formylphenyl)-2-methoxyiminoacetamide in 7 ml of methanol, was added 0.42 g (3.0 mmol) of α -methyl-4(trifluoromethyl)benzyloxyamine and the mixture was allowed to stand at room temperature overnight. After the reaction mixture was concentrated in vacuo, the

residue was chromatographed over SiO₂ to yield 0.89 g of the title compound (yield: 95 %).

The compounds Nos. 16-18, 20, 21, 23, 25, 27-29, 31, 32, 35, 38-40, 42, 44, 46, 47, 49-68 and 72 in Table 1 and the compounds Nos. 76, 77 and 79-83 in Table 2 were synthesized by repeating the above-mentioned procedure except altering the starting material.

5

SYNTHESIS EXAMPLE 4

Synthesis of methyl 2-[2-{4-(3-trifluoromethylphenyl)-2,3-diaza-1,3-pentadienyl}phenyl]-2-methoxyiminoacetate (compound No. 88 in Table 3):

10

A mixture comprising 1.5 g (6.75 mmol) of methyl 2-(2-formylphenyl)-2-methoxyiminoacetate, 1.37 g (6.75 mmol) of m-trifluoromethylacetophenone hydrazone and 7.5 ml of ethanol was heated under reflux for 3 hours. After vacuum concentration, the residue was chromatographed over SiO₂ to give 2.48 g of the title compound (yield: 86.6 %).

15

The compounds No. 86, No. 87, No. 89 and No. 90 in Table 3 were synthesized by repeating the above-mentioned procedure except altering the starting material.

SYNTHESIS EXAMPLE 5

20 Synthesis of methyl 2-[2-{4-(trifluoromethyl)benzoyloxyiminomethyl}phenyl]-2-methoxyiminoacetate (compound No. 91 in Table 3):

To a mixture comprising 0.60 g (2.53 mmol) of methyl 2-{2-(hydroxyiminomethyl)phenyl}-2-methoxyiminoacetate, 1 ml of triethylamine and 5 ml of dichloromethane, was added 0.80 g (3.8 mmol) of p-trifluoromethylbenzoic acid chloride under ice cooling. After stirring at room temperature for 12 hours, the reaction mixture was poured into water and extracted with ethyl acetate. Than it was successively washed with water and a saturated sodium chloride solution and dried over anhydrous sodium sulfate. After vacuum concentration, the residue was chromatographed over SiO₂ to give 0.31 g of the title compound (yield: 30.0 %).

30

35


40

45

50

55

Table 1



Compound No.	R ¹	R ²	X	(Y) _n	Property
1	H	H	H	3-CF ₃	m.p. 93-97°C
2	H	H	H	4-CF ₃	nD 1.5505/25°C
3	H	H	H	3-Cl	m.p. 90-93°C
4	H	H	H	4-Cl	m.p. 119-123°C
5	H	H	H	3-CH ₃	m.p. 90-93°C
6	H	H	H	4-CH ₃	m.p. 148-150°C
7	H	H	H	2,5-(CH ₃) ₂	m.p. 106.5-107°C
8	H	H	H	3-OCH ₃	viscous
9	H	H	H	4-OCH ₃	viscous
10	H	H	H	3-OCF ₃	viscous
11	H	H	H	4-OCF ₃	m.p. 96-97°C
12	H	H	H	2,5-Cl ₂	m.p. 148.5-150.5°C
13	H	H	H	3,5-(CF ₃) ₂	m.p. 102.5-105°C
14	CH ₃	H	H	-	m.p. 107.5-110.2°C
15	CH ₃	H	H	2-Cl	viscous
16	CH ₃	H	H	3-Cl	m.p. 82.5-83°C
17	CH ₃	H	H	4-Cl	m.p. 87.5-90.5°C
18	CH ₃	H	H	4-Cl	viscous
19	CH ₃	H	H	3-Br	viscous
20	CH ₃	H	H	4-Br	amorphous solid
21	CH ₃	H	H	3,4-Cl ₂	m.p. 110.9-111.7°C

Table 1 (continued)

5	Compound No.	R ¹	R ²	X	(Y) _n	Property
	22	CH ₃	H	H	2,5-Cl ₂	viscous
10	23	CH ₃	H	H	3-CF ₃	m.p. 88.5-89°C
	24	CH ₃	H	H	4-CF ₃	m.p. 70.5-71.5°C
	25	CH ₃	H	H	4-CF ₃	viscous
15	26	CH ₃	H	H	2-CH ₃	viscous
	27	CH ₃	H	H	3-CH ₃	viscous
20	28	CH ₃	H	H	4-CH ₃	viscous
	29	CH ₃	H	H	2,4-(CH ₃) ₂	viscous
	30	CH ₃	H	H	2,5-(CH ₃) ₂	viscous
25	31	CH ₃	H	H	3,4-(CH ₃) ₂	viscous
	32	CH ₃	H	H	4-C ₂ H ₅	amorphous solid
	33	CH ₃	H	H	4-C ₃ H ₇ (n)	viscous
30	34	CH ₃	H	H	4-C ₃ H ₇ (iso)	viscous
	35	CH ₃	H	H	4-C ₄ H ₉ (tert)	viscous
	36	CH ₃	H	H	2-OCH ₃	viscous
35	37	CH ₃	H	H	3-OCH ₃	viscous
	38	CH ₃	H	H	4-OCH ₃	amorphous solid
40	39	CH ₃	H	H	4-OC ₂ H ₅	amorphous solid
	40	CH ₃	H	H	4-OC ₃ H ₇ (n)	amorphous solid
	41	CH ₃	H	H	3-OC ₃ H ₇ (iso)	amorphous solid
45	42	CH ₃	H	H	4-OC ₃ H ₇ (iso)	amorphous solid
	43	CH ₃	H	H	3-propargyloxy	amorphous solid
	44	CH ₃	H	H	4-propargyloxy	amorphous solid
50	45	CH ₃	H	H	4-OCH ₂ CH=CCl ₂	viscous

Table 1 (continued)

Compound No.	R ¹	R ²	X	(Y) _n	Property
46	CH ₃	H	H	3-OPh	amorphous solid
47	CH ₃	H	H	4-OPh	amorphous solid
48	CH ₃	H	H	4-cyclopropyl-methyloxy	viscous
49	CH ₃	H	H	3-OCF ₃	m.p. 82-83°C
50	CH ₃	H	H	4-OCF ₃	amorphous solid
51	CH ₃	H	H	3-OCHF ₂	viscous
52	CH ₃	H	H	4-OCHF ₂	viscous
53	CH ₃	H	H	3-OCH ₂ CF ₃	viscous
54	CH ₃	H	H	4-OCH ₂ CF ₃	viscous
55	CH ₃	H	H	4-OCH ₂ CF ₂ CF ₃	viscous
56	CH ₃	H	H	3-OSO ₂ CF ₃	viscous
57	CH ₃	H	H	4-OSO ₂ CF ₃	viscous
58	CH ₃	H	H	3-OSO ₂ C ₂ H ₅	viscous
59	CH ₃	H	H	4-OSO ₂ C ₂ H ₅	viscous
60	CH ₃	H	H	4-NHCOCF ₃	viscous
61	CH ₃	H	H	4-CN	m.p. 122.8-123.7°C
62	CH ₃	H	H	4-CN	amorphous solid
63	CH ₃	H	H	4-Ph	m.p. 65-70.9°C
64	CH ₃	H	H	3-SCH ₃	viscous
65	CH ₃	H	H	4-SCH ₃	viscous
66	CH ₃	H	H	3-SC ₃ H ₇ (iso)	viscous
67	CH ₃	H	H	4-SC ₃ H ₇ (iso)	viscous
68	CN	H	H	4-CF ₃	viscous
69	C ₂ H ₅	CH ₃	H	3-Cl	viscous


Table 1 (continued)

Compound No.	R ¹	R ²	X	(Y) _n	Property
70	C ₂ H ₅	H	H	4-OCOC ₂ H ₅	viscous
71	C ₂ H ₅	H	H	4-OCF ₃	viscous
72	CN	H	H	-	viscous
73	CF ₃	H	H	4-CF ₃	viscous
74	CH ₃	H	3-Cl	-	viscous
75	CH ₃	H	3-Cl	4-CF ₃	viscous

* The compounds No. 18, No. 25 and No. 62 are respectively isomers of the compounds No. 17, No. 24 and No. 61 at the benzyloxyimino moiety.

** Although E- and Z-isomers at the methoxyimino moiety exist, the properties of E-isomers alone are given in the above table.


Table 2



Compound No.	R ¹	R ²	Ar	Property
76	CH ₃	H	2-naphthyl	amorphous solid
77	CH ₃	H	1-naphthyl	viscous
78	H	H	6-chloropyridine-2-yl	viscous
79	CH ₃	H	6-trifluoromethyl-pyridine-2-yl	viscous
80	CH ₃	H	3,4-methylenedioxyphenyl	amorphous solid
81	CH ₃	H	3,4-ethylenedioxyphenyl	viscous
82	CH ₃	H	5-chloro-2-thienyl	viscous
83	CH ₃	H	2-chloro-4-methylthiazole-5-yl	viscous
84	CH ₃	CH ₃	2,4-dimethylthiazole-5-yl	viscous
85	C ₂ H ₅	H	2-t-butylthiazole-5-yl	viscous

* Although E- and Z-isomers at the methoxyimino moiety exist, the properties of the E-isomers alone are given in the above table.

Table 3



Compound No.	B	(Y) _n	Property
86	N=C(CH ₃)	3-CH ₃	97-104°C
87	N=C(CH ₃)	3-Cl	117-118.5°C
88	N=C(CH ₃)	3-CF ₃	138-138.5°C
89	N=C(CH ₃)	4-Cl	129-132°C
90	N=C(CH ₃)	4-CF ₃	141-144°C
91	O-C(O)	4-CF ₃	nD 1.5474/25

* Although E- and Z-isomers at the methoxyimino moiety exist, the properties of the E-isomers alone are given in the above table.

¹H-NMR data of the obtained compounds are as follows.

[Table 1]

¹H-NMR data of compound No. 1 (CDCl₃); 2.89 (3H, d), 3.89 (3H, s), 5.20 (2H, s), 6.73 (1H, brs), 7.18 (1H, dd), 7.39 (1H, dd), 7.42 (1H, d), 7.49 (1H, d), 7.55-7.60 (2H, m), 7.64 (1H, s), 7.76 (1H, dd), 7.97 (1H, s)

¹H-NMR data of compound No. 2 (CDCl₃); 2.79 (3H, d), 3.85 (3H, s), 5.18 (2H, s), 6.95 (1H, br), 7.19 (1H, d), 7.32-7.40 (2H, m), 7.47 (2H, d), 7.59 (2H, d), 7.76 (1H, d), 8.03 (1H, s)

¹H-NMR data of compound No. 3 (CDCl₃); 2.89 (3H, d), 3.90 (3H, s), 5.11 (2H, s), 6.71 (1H, brs), 7.18 (1H, dd), 7.24-7.30 (3H, m), 7.37 (1H, s), 7.39 (1H, dd), 7.41 (1H, dd), 7.74 (1H, dd), 7.96 (1H, s)

¹H-NMR data of compound No. 4 (CDCl₃); 2.89 (3H, d), 3.90 (3H, s), 5.11 (2H, s), 6.69 (1H, brs), 7.17 (1H, dd), 7.32 (4H, s), 7.39 (1H, dd), 7.41 (1H, dd), 7.75 (1H, dd), 7.94 (1H, s)

¹H-NMR data of compound No. 5 (CDCl₃); 2.35 (3H, s), 2.83 (3H, d), 3.87 (3H, s), 5.11 (2H, s), 6.71 (1H, brs), 7.11 (1H, d), 7.15-7.27 (4H, m), 7.36 (1H, dd), 7.39 (1H, dd), 7.76 (1H, dd), 7.95 (1H, s)

¹H-NMR data of compound No. 6 (CDCl₃); 2.35 (3H, s), 2.88 (3H, d), 3.90 (3H, s), 5.11 (2H, s), 6.67 (1H, brs), 7.17 (2H, d), 7.18 (1H, dd), 7.28 (2H, d), 7.38 (1H, dd), 7.41 (1H, dd), 7.77 (1H, dd), 7.93 (1H, s)

¹H-NMR data of compound No. 7 (CDCl₃); 2.33 (3H, s), 2.34 (3H, s), 2.88 (3H, d), 3.91 (3H, s), 5.15 (2H, s), 6.68 (1H, brs), 7.05 (1H, d), 7.08 (1H, d), 7.16 (1H, s), 7.18 (1H, dd), 7.39 (1H, dd), 7.42 (1H, dd), 7.80 (1H, dd), 7.94 (1H, s)

¹H-NMR data of compound No. 8 (CDCl₃); 2.86 (3H, s), 3.81 (3H, s), 3.90 (3H, s), 5.15 (2H, s), 6.76 (1H, br), 6.87 (1H, d), 6.96 (1H, s), 6.98 (1H, d), 7.20 (1H, dd), 7.29 (1H, dd), 7.39 (1H, dd), 7.42 (1H, dd), 7.78 (1H, dd), 7.99 (1H, s)

¹H-NMR data of compound No. 11 (CDCl₃); 2.88 (3H, d), 3.89 (3H, s), 5.14 (2H, s), 6.71 (1H, br), 7.16-7.24 (3H, m), 7.38-7.44 (4H, m), 7.76 (1H, dd), 7.95 (1H, s)

¹H-NMR data of compound No. 12 (CDCl₃); 2.90 (3H, d), 3.92 (3H, s), 5.23 (2H, s), 6.75 (1H, brs), 7.19 (1H,

- dd), 7.22 (1H, dd), 7.30 (1H, d), 7.39-7.45 (3H, m), 7.75 (1H, dd), 8.01 (1H, s)
¹H-NMR data of compound No. 13 (CDCl₃); 2.91 (3H, d), 3.89 (3H, s), 5.24 (2H, s), 6.77 (1H, br), 7.18 (1H, dd), 7.37-7.46 (2H, m), 7.76 (1H, dd), 7.83 (3H, s), 7.98 (1H, s)
¹H-NMR data of compound No. 14 (CDCl₃); 1.58 (3H, d), 2.86 (3H, d), 3.87 (3H, s), 5.26 (1H, g), 6.56 (1H, br), 7.16 (1H, m), 7.3-7.4 (7H), 7.68 (1H, m), 7.96 (1H, s)
5 ¹H-NMR data of compound No. 16 (CDCl₃); 1.55 (3H, d), 2.87 (3H, d), 3.88 (3H, s), 5.22 (1H, q), 6.63 (1H, br), 7.15 (1H, dd), 7.19-7.28 (3H, m), 7.33 (1H, s), 7.36 (1H, dd), 7.39 (1H, dd), 7.67 (1H, dd), 7.96 (1H, s)
¹H-NMR data of compound No. 17 (CDCl₃); 1.56 (3H, d), 2.83 (3H, d), 3.87 (3H, s), 5.26 (1H, q), 6.76 (1H, brs), 7.17 (1H, dd), 7.26-7.40 (6H, m), 7.70 (1H, dd), 7.98 (1H, s)
10 ¹H-NMR data of compound No. 18 (CDCl₃); 1.55 (3H, d), 2.90 (3H, d), 3.90 (3H, s), 5.25 (1H, q), 6.77 (1H, br), 7.2-8.3 (9H, m)
¹H-NMR data of compound No. 20 (CDCl₃); 1.55 (3H, d), 2.88 (3H, d), 3.88 (3H, s), 5.22 (1H, q), 6.6 (1H, br), 7.16 (1H, m), 7.22 (2H, d), 7.38 (2H, m), 7.47 (2H, d), 7.68 (1H, m), 7.94 (1H, s)
¹H-NMR data of compound No. 21 (CDCl₃); 1.54 (3H, d), 2.89 (3H, d), 3.89 (3H, s), 5.20 (1H, q), 6.68 (1H, br), 7.18 (2H, m), 7.4 (4H, m), 7.68 (1H, m), 7.95 (1H, s)
15 ¹H-NMR data of compound No. 23 (CDCl₃); 1.58 (3H, d), 2.88 (3H, d), 3.86 (3H, s), 5.30 (1H, g), 6.67 (1H, br), 7.15 (1H, dd), 7.35-7.55 (5H, m), 7.60 (1H, s), 7.68 (1H, dd), 7.96 (1H, s)
¹H-NMR data of compound No. 24 (CDCl₃); 1.57 (3H, d), 2.87 (3H, d), 3.86 (3H, s), 5.31 (1H, q), 6.63 (1H, br), 7.15 (1H, dd), 7.35-7.40 (2H, m), 7.46 (2H, d), 7.60 (2H, d), 7.67 (1H, dd), 7.97 (1H, s)
20 ¹H-NMR data of compound No. 25 (CDCl₃); 1.56 (3H, d), 2.87 (3H, d), 3.88 (3H, s), 5.32 (1H, q), 6.82 (1H, br), 7.2-8.3 (9H, m)
¹H-NMR data of compound No. 27 (CDCl₃); 1.59 (3H, d), 2.38 (3H, s), 2.85 (3H, d), 3.89 (3H, s), 5.26 (1H, q), 6.63 (1H, br), 7.08-7.30 (5H, m), 7.37 (1H, dd), 7.40 (1H, dd), 7.71 (1H, dd), 7.98 (1H, s)
¹H-NMR data of compound No. 28 (CDCl₃); 1.56 (3H, d), 2.34 (3H, s), 2.86 (3H, d), 3.88 (3H, s), 5.23 (1H, q), 6.58 (1H, br), 7.15 (1H, dd), 7.16 (2H, d), 7.24 (2H, d), 7.36 (1H, dd), 7.39 (1H, dd), 7.68 (1H, dd), 7.94 (1H, s)
25 ¹H-NMR data of compound No. 29 (CDCl₃); 1.55 (3H, d), 2.30 (3H, s), 2.35 (3H, s), 2.86 (3H, d), 3.89 (3H, s), 5.47 (1H, q), 6.55 (1H, brs), 6.99 (1H, d), 7.02 (1H, d), 7.17 (1H, dd), 7.27 (1H, dd), 7.37 (2H, m), 7.70 (1H, dd), 7.94 (1H, s)
¹H-NMR data of compound No. 31 (CDCl₃); 1.56 (3H, d), 2.25 (3H, s), 2.27 (3H, s), 2.86 (3H, d), 3.88 (3H, s), 5.20 (1H, q), 6.60 (1H, brs), 7.1-7.2 (4H, m), 7.37 (2H, m), 7.70 (1H, dd), 7.94 (1H, s)
¹H-NMR data of compound No. 32 (CDCl₃); 1.24 (3H, t), 1.57 (3H, d), 2.64 (2H, q), 2.86 (3H, d), 3.87 (3H, s), 5.23 (1H, q), 6.56 (1H, br), 7.17 (1H, m), 7.18 (2H, d), 7.27 (2H, d), 7.37 (2H, m), 7.7 (1H, m), 7.95 (1H, s)
35 ¹H-NMR data of compound No. 35 (CDCl₃); 1.28 (9H, s), 1.58 (3H, d), 2.86 (3H, d), 3.87 (3H, s), 5.27 (1H, q), 6.6 (1H, br), 7.16 (1H, m), 7.29 (2H, d), 7.38 (4H, m), 7.7 (1H, m), 7.95 (1H, s)
¹H-NMR data of compound No. 38 (CDCl₃); 1.57 (3H, d), 2.87 (3H, d), 3.81 (3H, s), 3.88 (3H, s), 5.22 (1H, q), 6.62 (1H, br), 6.90 (2H, d), 7.16 (1H, m), 7.30 (2H, d), 7.38 (2H, m), 7.69 (1H, m), 7.93 (1H, s)
¹H-NMR data of compound No. 39 (CDCl₃); 1.41 (3H, t), 1.56 (3H, d), 2.87 (3H, d), 3.89 (3H, s), 4.02 (2H, q), 6.6 (1H, br), 6.88 (2H, d), 7.16 (1H, m), 7.3 (2H, d), 7.38 (2H, m), 7.7 (1H, m), 7.93 (1H, s)
40 ¹H-NMR data of compound No. 40 (CDCl₃); 1.03 (3H, t), 1.57 (3H, d), 1.8 (2H, q), 2.87 (3H, d), 3.89 (3H, s), 3.91 (2H, t), 5.21 (1H, q), 6.6 (1H, br), 6.9 (2H, d), 7.18 (1H, m), 7.3 (2H, d), 7.37 (2H, m), 7.7 (1H, m), 7.93 (1H, s)
¹H-NMR data of compound No. 42 (CDCl₃); 1.33 (6H, d), 1.56 (3H, d), 2.87 (3H, d), 3.89 (3H, d), 4.52 (1H, m), 5.21 (1H, g), 6.62 (1H, br), 6.85 (2H, d), 7.16 (1H, m), 7.27 (2H, d), 7.38 (2H, m), 7.70 (1H, m), 7.93 (1H, s)
45 ¹H-NMR data of compound No. 44 (CDCl₃); 1.57 (3H, d), 2.50 (1H, t), 2.87 (3H, d), 3.88 (3H, s), 4.69 (2H, d), 5.21 (1H, q), 6.61 (1H, br), 6.97 (2H, d), 7.16 (1H, m), 7.30 (2H, d), 7.38 (2H, m), 7.70 (1H, m), 7.93 (1H, s)
¹H-NMR data of compound No. 49 (CDCl₃); 1.56 (3H, d), 2.88 (3H, d), 3.87 (3H, s), 5.57 (1H, q), 6.67 (1H, br), 7.13 (1H, d), 7.15 (1H, dd), 7.20 (1H, s), 7.28 (1H, d), 7.34-7.42 (3H, m), 7.68 (1H, dd), 7.96 (1H, s)
¹H-NMR data of compound No. 50 (CDCl₃); 1.57 (3H, d), 2.88 (3H, d), 3.87 (3H, s), 5.28 (1H, q), 6.64 (1H, br), 7.16 (2H, dd), 7.19 (2H, d), 7.38 (4H, m), 7.68 (1H, dd), 7.95 (1H, s)
¹H-NMR data of compound No. 54 (CDCl₃); 1.56 (3H, d), 2.88 (3H, d), 3.89 (3H, s), 4.35 (2H, q), 5.25 (1H, q), 6.64 (1H, br), 6.93 (2H, d), 7.17 (1H, dd), 7.32 (2H, d), 7.38 (2H, m), 7.70 (1H, dd), 7.93 (1H, s)
55 ¹H-NMR data of compound No. 55 (CDCl₃); 1.57 (3H, d), 2.89 (3H, d), 3.89 (3H, s), 4.42 (2H, t), 5.23 (1H, q), 6.68 (1H, br), 6.93 (2H, d), 7.16 (1H, dd), 7.32 (2H, d), 7.38 (2H, m), 7.7 (1H, dd), 7.93 (1H, s)
¹H-NMR data of compound No. 56 (CDCl₃); 1.56 (3H, d), 2.86 (3H, d), 3.86 (3H, s), 5.29 (1H, q), 6.80 (1H,

br), 7.14-7.21 (2H, m), 7.27 (1H, s), 7.33-7.45 (4H, m), 7.68 (1H, dd), 7.98 (1H, s)

¹H-NMR data of compound No. 58 (CDCl₃): 1.47 (3H, t), 1.56 (3H, d), 2.86 (3H, d), 3.24 (2H, q), 3.84 (3H, s), 5.27 (1H, q), 6.77 (1H, br), 7.14 (1H, dd), 7.18 (1H, d), 7.25 (1H, s), 7.29 (1H, d), 7.33-7.41 (3H, m), 7.66 (1H, dd), 7.96 (1H, s)

5 ¹H-NMR data of compound No. 61 (CDCl₃): 1.56 (3H, d), 2.89 (3H, d), 3.89 (3H, s), 5.30 (1H, g), 6.66 (1H, br), 7.15 (1H, m), 7.38 (2H, m), 7.42 (2H, d), 7.64 (2H, d), 7.68 (1H, m), 7.96 (1H, m)

¹H-NMR data of compound No. 62 (CDCl₃): 1.54 (3H, d), 2.87 (3H, d), 3.87 (3H, s), 5.22 (1H, q), 6.64 (1H, br), 7.16 (1H, m), 7.22 (2H, d), 7.37 (2H, m), 7.47 (2H, d), 7.66 (1H, m), 7.94 (1H, s)

10 ¹H-NMR data of compound No. 63 (CDCl₃): 1.62 (3H, d), 2.85 (3H, d), 3.87 (3H, s), 5.34 (1H, q), 6.60 (1H, br), 7.18 (1H, m), 7.3-7.7 (12H), 7.98 (1H, s)

[Table 2]

15 ¹H-NMR data of compound No. 76 (CDCl₃): 1.66 (3H, d), 2.78 (3H, d), 3.81 (3H, s), 5.42 (1H, q), 6.5 (1H, br), 7.15 (1H, m), 7.36 (2H, m), 7.5 (3H, m), 7.66 (1H, m), 7.8 (4H, m), 8.01 (1H, s)

¹H-NMR data of compound No. 78 (CDCl₃): 2.88 (3H, d), 3.88 (3H, s), 5.23 (2H, s), 6.82 (1H, br), 7.18 (1H, dd), 7.24 (1H, d), 7.33 (1H, d), 7.39 (1H, dd), 7.42 (1H, dd), 7.66 (1H, dd), 7.70 (1H, dd), 8.04 (1H, s)

20 ¹H-NMR data of compound No. 80 (CDCl₃): 1.54 (3H, d), 2.89 (3H, d), 3.89 (3H, s), 5.18 (1H, q), 5.94 (2H, s), 6.65 (1H, br), 6.7-6.9 (3H, m), 7.16 (1H, m), 7.38 (2H, m), 7.69 (1H, m), 7.93 (1H, s)

¹H-NMR data of compound No. 81 (CDCl₃): 1.54 (3H, d), 2.89 (3H, d), 3.89 (3H, s), 4.25 (4H, s), 5.15 (1H, q), 6.66 (1H, br), 6.84 (2H, d), 6.88 (1H, d), 7.15 (1H, m), 7.38 (2H, m), 7.70 (1H, m), 7.92 (1H, s)

¹H-NMR data of compound No. 82 (CDCl₃): 1.64 (3H, d), 2.91 (3H, d), 3.92 (3H, s), 5.36 (1H, g), 6.75 (1H, br), 6.78 (2H, m), 7.19 (1H, m), 7.4 (2H, m), 7.75 (1H, m), 7.90 (1H, s)

25

[Table 3]

¹H-NMR data of compound No. 86 (CDCl₃): 2.41 (3H, s), 2.46 (3H, s), 3.82 (3H, s), 4.03 (3H, s), 7.23-7.35 (3H, m), 7.48-7.54 (2H, m), 7.66 (1H, d), 7.73 (1H, s), 7.87 (1H, dd), 8.37 (1H, s)

30 ¹H-NMR data of compound No. 87 (CDCl₃): 2.44 (3H, s), 3.81 (3H, s), 4.03 (3H, s), 7.26 (1H, dd), 7.31-7.43 (2H, m), 7.48-7.54 (2H, m), 7.76 (1H, d), 7.86 (1H, dd), 7.91 (1H, s), 8.37 (1H, s)

¹H-NMR data of compound No. 88 (CDCl₃): 2.49 (3H, s), 3.82 (3H, s), 4.03 (3H, s), 7.27 (1H, dd), 7.51 (2H, dd), 7.54 (1H, d), 7.67 (1H, d), 7.87 (1H, dd), 8.06 (1H, d), 8.19 (1H, s), 8.39 (1H, s)

35 ¹H-NMR data of compound No. 91 (CDCl₃): 3.92 (3H, s), 4.06 (3H, s), 7.29 (1H, d), 7.49-7.62 (2H, m), 7.77 (2H, d), 8.10 (1H, d), 8.23 (2H, d), 8.38 (1H, s)

FORMULATION EXAMPLE 1

40 A wettable powder was obtained by uniformly pulverizing and mixing 20 parts of the compound No. 2 given in Table 1, 75 parts of diatomaceous earth and 5 parts of a surfactant comprising alkyl benzenesulfonate as a main component.

FORMULATION EXAMPLE 2

45 An emulsifiable concentrate was obtained by mixing and dissolving 30 parts of the compound No. 3 given in Table 1, 15 parts of "Sorpel®" 3005X (a nonionic surfactant/anionic surfactant mixture manufactured by Toho Chemical Industry Co., Ltd.), 25 parts of xylene and 30 parts of dimethylformamide.

To clarify the usefulness of the compounds of the present invention as an agricultural/horticultural fungicide, the following Test Examples will be given.

50

TEST EXAMPLE 1

Preventive activity on wheat powdery mildew

55 A wettable powder prepared in the same manner as described in Formulation Example 1 was diluted with water to a definite concentration and then applied by foliar application on wheat plants (var. Norin No. 61) at the 1 to 2 leaf stage grown in pots of 6 cm in diameter at a ratio of 10 ml/pot. After air-drying the chemical solution, a spore suspension of *Erysiphe graminis* (a pathogen of wheat powdery mildew) was

EP 0 596 254 A1

inoculated to the plants by spraying. Then the plants were kept in a greenhouse for 7 to 10 days.

For evaluation, the diseased area ratio of each leaf was measured and the preventive value was calculated in accordance with the following formula. The results are listed as "Preventive value 1" in Table 4.

5

$$\begin{array}{l} \text{(average diseased area ratio} \\ \text{in untreated plot) - (average} \\ \text{diseased area ratio in} \\ \text{treated plot)} \\ \text{Preventive value (\%)} = \frac{\quad}{\text{(average diseased area ratio} \\ \text{in untreated plot)}} \times 100 \end{array}$$

15

The compound numbers correspond to the compound Nos. in Tables 1, 2 and 3.

TEST EXAMPLE 2

Preventive activity on wheat brown rust

20

A wettable powder prepared in the same manner as described in Test Example 1 was diluted with water to a definite concentration and then applied by foliar application on wheat plants (var. Norin No. 61) at the 1 to 2 leaf stage grown in pots of 6 cm in diameter at a ratio of 10 ml/pot. After air-drying the chemical solution, a spore suspension of *Puccinia recondita* (a pathogen of wheat brown rust) was inoculated into the plants by spraying. Then the plants were kept in a moist chamber at 22 °C for 15 hours and then allowed to stand in a greenhouse for 7 days.

25

For evaluation, the diseased area ratio of each leaf was measured and the preventive value was calculated in accordance with the following formula. The results are listed as "Preventive value 2" in Table 4.

30

$$\begin{array}{l} \text{(average diseased area ratio} \\ \text{in untreated plot) - (average} \\ \text{diseased area ratio in} \\ \text{treated plot)} \\ \text{Preventive value (\%)} = \frac{\quad}{\text{(average diseased area ratio} \\ \text{in untreated plot)}} \times 100 \end{array}$$

35

40

The compound numbers correspond to the compound Nos. in Tables 1, 2 and 3.

45

50

55

Table 4

Compound No.	Active ingredient (ppm)	Preventive value 1 (%)	Preventive value 2 (%)
1	200	100	100
2	200	100	100
3	200	100	100
4	200	98	99
5	200	99	100
6	200	98	96
7	200	85	99
8	200	99	99
9	200	95	98
10	200	99	100
11	200	100	99
12	200	85	80
13	200	99	98
14	200	100	100
15	200	93	96
16	200	100	99
17	200	100	100
18	200	100	100
19	200	100	100
20	200	100	100
21	200	100	100
22	200	98	90
23	200	100	99
24	200	100	100
25	200	100	100
26	200	89	93

Table 4 (continued)

5	Compound No.	Active ingredient (ppm)	Preventive value 1 (%)	Preventive value 2 (%)
	27	200	100	100
10	28	200	100	100
	29	200	100	98
	30	200	99	97
15	31	200	100	100
	32	200	100	100
	33	200	100	100
20	34	200	100	100
	35	200	100	100
	36	200	98	98
25	37	200	100	100
	38	200	100	100
	39	200	100	100
30	40	200	100	100
	41	200	100	100
	42	200	100	100
35	43	200	100	100
	44	200	100	100
	45	200	100	100
40	46	200	100	100
	47	200	100	100
	48	200	100	100
45	49	200	100	100
	50	200	100	100
	51	200	100	100
50	52	200	100	100

Table 4 (continued)

5	Compound No.	Active ingredient (ppm)	Preventive value 1 (%)	Preventive value 2 (%)
	53	200	100	100
10	54	200	100	100
	55	200	100	100
	56	200	100	100
15	57	200	100	100
	58	200	100	100
	59	200	100	100
20	60	200	100	100
	61	200	100	100
	62	200	100	100
25	63	200	100	100
	64	200	100	100
	65	200	100	100
30	66	200	100	100
	67	200	100	100
	68	200	95	92
35	69	200	89	99
	70	200	99	95
	71	200	100	100
40	72	200	95	92
	73	200	98	95
	74	200	100	100
45	75	200	100	100
	76	200	100	100
	77	200	100	100
50	78	200	90	87

Table 4 (continued)

Compound No.	Active ingredient (ppm)	Preventive value 1 (%)	Preventive value 2 (%)
79	200	100	100
80	200	100	100
81	200	100	100
82	200	99	98
83	200	100	100
84	200	90	89
85	200	98	96

Residual activities were evaluated in cases simulating the practical uses as shown in following Test Examples. The tests were performed on the compounds according to the present invention as well as ones described in the prior arts, in order to clarify the advancement of the present invention over prior arts.

TEST EXAMPLE 3

Residual activity on wheat powdery mildew

A wettable powder prepared in the same manner as described in Formulation Example 1 was diluted with water to a definite concentration and then applied by foliar application to wheat plants (var. Norin No. 61) at the 1 to 2 leaf stage grown in pots of 6 cm in diameter at a ratio of 10 ml/pot. After keeping outdoor for 14 days, the plants were inoculated with a spore suspension of *Erysiphe graminis* (a pathogen of wheat powdery mildew) by spraying. Then the plants were allowed to stand in a greenhouse at 20 to 22 °C for 7 to 10 days.

For evaluation, the diseased area ratio of each leaf was measured and the preventive value was calculated in accordance with the following formula. The results are listed as "Preventive value 3" in Table 5.

$$\text{Preventive value (\%)} = \frac{(\text{average diseased area ratio in untreated plot}) - (\text{average diseased area ratio in treated plot})}{(\text{average diseased area ratio in untreated plot})} \times 100$$

The compound numbers correspond to the compound Nos. in Tables 1, 2 and 3.

TEST EXAMPLE 4

Residual activity on wheat brown rust

A wettable powder prepared in the same manner as described in Test Example 1 was diluted with water to a definite concentration and then applied by foliar application to wheat plants (var. Norin No. 61) at the 1 to 2 leaf stage grown in pots of 6 cm in diameter at a ratio of 10 ml/pot. After keeping outdoor for 14 days,

the plants were inoculated with a spore suspension of *Puccinia recondita* (a pathogen of wheat brown rust) by spraying. Then the plants were kept in a moist chamber at 22 °C for 24 hours and then allowed to stand in a greenhouse at 20 to 25 °C for 10 days.

For evaluation, the diseased area ratio of each leaf was measured and the preventive value was calculated in accordance with the following formula. The results are listed as "Preventive value 4" in Table 5.

$$\text{Preventive value (\%)} = \frac{(\text{average diseased area ratio in untreated plot}) - (\text{average diseased area ratio in treated plot})}{(\text{average diseased area ratio in untreated plot})} \times 100$$

The compound numbers correspond to the compound Nos. in Tables 1, 2 and 3.

Table 5








U	V		 250 50 10 (ppm)	 250 50 10 (ppm)	 250 50 10 (ppm)
H	3-CF ₃ -Phenyl	Preventive value 3 Preventive value 4 referential patent No.	100 62 12 100 77 23 invention compound No. 1	86 0 0 68 0 0 EP499823	70 0 0 98 13 0 WO92/13830
CH ₃	Phenyl	Preventive value 3 Preventive value 4 referential patent No.	100 88 59 100 85 11 invention compound No. 14	99 48 0 99 36 0 EP499823	99 55 0 98 38 0 EP463488
CH ₃	4-Cl-Phenyl	Preventive value 3 Preventive value 4 referential patent No.	100 92 33 100 97 27 invention compound No. 17	100 54 0 95 31 0 EP499823	100 78 0 100 83 0 EP463488
CH ₃	4-CH ₃ -Phenyl	Preventive value 3 Preventive value 4 referential patent No.	100 82 23 100 58 18 invention compound No. 28	88 0 0 53 0 0 EP499823	99 50 0 98 45 0 EP463488

Table 5 (continued)

U	V			250 50 10 (ppm)		250 50 10 (ppm)		250 50 10 (ppm)
CH ₃	4-CF ₃ - Phenyl	Preventive value 3 Preventive value 4 referential patent No.	100 99 96 100 96 91 invention compound No. 24	100 99 96 100 96 91 invention compound No. 24	99 0 0 99 24 0 EP499823	99 0 0 99 24 0 EP499823	100 70 15 99 77 28 EP463488	
CH ₃	2-Naphthyl	Preventive value 3 Preventive value 4 referential patent No.	100 99 95 100 95 92 invention compound No. 76	100 99 95 100 95 92 invention compound No. 76	85 20 0 92 48 0 EP499823	85 20 0 92 48 0 EP499823	92 77 0 90 0 0 EP463488	
						250 50 10 (ppm)		
		Preventive value 3 Preventive value 4 referential patent No.			52 8 0 16 0 0 EP398692			

Each of the compounds of the present invention is a novel one having an excellent fungicidal activity. It is particularly effective in controlling phytopathogenic fungi, which makes it highly useful as an agricultural/horticultural fungicide.


For example, these compounds exert high activity on rice blast (*Pyricularia oryzae*), rice sheath blight (*Rhizoctonia solani*), wheat powdery mildew (*Erysiphe graminis f. sp. tritici*) and barley powdery mildew (*E. graminis f. sp. hodei*), various leaf rusts of wheat and barley (e.g., *Puccinia recondita*), gray mold of vegetables and fruit trees (*Botrytis cinerea*) and late blight of various crops (*Phytophthora infestans*).

Further, they have prolonged residual activity and excellent systemic action in plants, which makes them useful as an agricultural/horticultural fungicide.

While the invention has been described in detail and with reference to specific embodiments thereof, it will be apparent to one skilled in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof.


Claims

1. A methoxyiminoacetic acid derivative represented by the following formula (I):



wherein X represents a hydrogen atom, a halogen atom, an alkyl group having 1 to 4 carbon atoms or an alkoxy group having 1 to 4 carbon atoms; A represents a methoxy group or a methylamino group; when A is a methoxy group, B represents -O-CO- or -N=C(R¹)- and when A is a methylamino group, B represents -O-CR¹R²-, wherein R¹ and R² independently represent a hydrogen atom, an alkyl group having 1 to 4 carbon atoms, a cyano group or a trifluoromethyl group; and Ar represents an optionally substituted aryl group or an optionally substituted heteroaryl group.

2. A methoxyiminoacetic acid derivative represented by the following formula (II):



wherein X represents a hydrogen atom or a halogen atom; R¹ and R² independently represent a hydrogen atom, an alkyl group having 1 to 4 carbon atoms, a cyano group or a trifluoromethyl group; and Ar represents an optionally substituted aryl group or an optionally substituted heteroaryl group.

3. The methoxyiminoacetic acid derivative as claimed in Claim 1, wherein Ar represents

an optionally substituted aryl group by 1 to 5 substituents which may be the same or different and, when the aryl group have 2 or more substituents, those adjacent to each other may be combined together to give a methylenedioxy or ethylenedioxy group and form a fused ring together with the aryl group, said substituents being selected from the group consisting of a cyano group; a halogen atom; an alkyl group having 1 to 6 carbon atoms; an alkenyl group having 2 to 4 carbon atoms optionally substituted by a halogen atom; a haloalkyl group having 1 to 4 carbon atoms; an alkoxy group having 1 to 6 carbon atoms optionally substituted by a halogen atom or a cycloalkyl group having 3 to 6 carbon atoms; an alkylcarbonyloxy group having 1 to 7 carbon atoms optionally substituted by a halogen atom; an acylamino group having 1 to 7 carbon atoms optionally substituted by a halogen atom; an alkylthio group having 1 to 6 carbon atoms optionally substituted by a halogen atom; an aryl group optionally substituted by an alkyl group having 1 to 4 carbon atoms or a halogen atom; an aryloxy group optionally substituted by an alkyl group having 1 to 4 carbon atoms or a halogen atom; an alkylsul-

fonyloxy group having 1 to 6 carbon atoms optionally substituted by a halogen atom; an alkenyloxy group having 2 to 6 carbon atoms optionally substituted by a halogen atom, and an alkynyloxy group having 2 to 6 carbon atoms; or

an optionally substituted heteroaryl group by 1 to 2 substituents which may be the same or different, said substituents being selected from the group consisting of a cyano group; a halogen atom; an alkyl group having 1 to 6 carbon atoms; a haloalkyl group having 1 to 4 carbon atoms; and an alkoxy group having 1 to 6 carbon atoms optionally substituted by a halogen atom; or a cycloalkyl group having 3 to 6 carbon atoms.

4. The methoxyiminoacetic acid derivative as claimed in Claim 2, wherein X represents a hydrogen atom or a halogen atom; R¹ and R² independently represent a hydrogen atom, a cyano group or an alkyl group having 1 to 4 carbon atoms; and Ar represents

an optionally substituted phenyl group or an optionally substituted naphthyl group, by one or more substituents being selected from the group consisting of a halogen atom; an alkyl group having 1 to 4 carbon atoms; an alkoxy group having 1 to 4 carbon atoms optionally substituted by a halogen atom; an acylamino group having 1 to 4 carbon atoms optionally substituted by a halogen atom; an alkylthio group having 1 to 3 carbon atoms; an alkylsulfonyloxy group having 1 to 3 carbon atoms optionally substituted by a halogen atom; or a trifluoromethyl group, or

an optionally substituted thienyl group or an optionally substituted thiazolyl group, by one or more substituents selected from the group consisting of a halogen atom; an alkyl group having 1 to 4 carbon atoms; or a trifluoromethyl group.

5. The methoxyiminoacetic acid derivative as claimed in Claim 2, wherein X represents a hydrogen atom; R¹ represents a methyl group or a cyano group; and R² represents a hydrogen atom; and Ar represents a naphthyl group or an optionally substituted phenyl group by one or more substituents being selected from the group consisting of a halogen atom; an alkyl group having 1 to 4 carbon atoms; an alkoxy group having 1 to 4 carbon atoms optionally substituted by a halogen atom; an acylamino group having 1 to 4 carbon atoms optionally substituted by a halogen atom; an alkylsulfonyloxy group having 1 to 3 carbon atoms optionally substituted by a halogen atom; or a trifluoromethyl group.

6. The methoxyiminoacetic acid derivative as claimed in Claim 5, wherein Ar represents an optionally substituted phenyl group by one or more substituents being selected from the group consisting of a halogen atom; an alkyl group having 1 to 4 carbon atoms; an alkoxy group having 1 to 4 carbon atoms optionally substituted by a fluorine; an alkylsulfonyloxy group having 1 to 3 carbon atoms optionally substituted by a fluorine; or a trifluoromethyl group.

7. An agricultural/horticultural fungicide containing a methoxyiminoacetic acid derivative as claimed in Claim 1 as an active ingredient.

8. An agricultural/horticultural fungicide containing a methoxyiminoacetic acid derivative as claimed in Claim 2 as an active ingredient.

9. An agricultural/horticultural fungicide containing a methoxyiminoacetic acid derivative as claimed in Claim 3 as an active ingredient.

10. An agricultural/horticultural fungicide containing a methoxyiminoacetic acid derivative as claimed in Claim 4 as an active ingredient.

11. An agricultural/horticultural fungicide containing a methoxyiminoacetic acid derivative as claimed in Claim 5 as an active ingredient.

12. An agricultural/horticultural fungicide containing a methoxyiminoacetic acid derivative as claimed in Claim 6 as an active ingredient.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 93 11 5917

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X,D	EP-A-0 499 823 (BASF) * compound I and page 3, line 33 *	1,3,7	C07C251/52 A01N37/50
Y,D	* table I, compound 1.24 - 1.234, 1.397 - 1.461, claims 1, 3, 4, 5 *	1-12	C07C251/48 C07C251/54 C07C251/58
Y,D	EP-A-0 398 692 (SHIONOGI SEIYAKU KABUSHIKI) * page 3, line 11 - line 41; claims 1,14-16 *	1-12	C07C255/62 C07C251/88 C07C251/68 C07D213/61 C07D317/54
A,D	WO-A-92 13830 (IMPERIAL CHEMICAL INDUSTRIES) * claims *	1-12	C07D333/28 C07D277/32
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C07C
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 6 January 1994	Examiner Seufert, G
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document	

EPF FORM 150 01.82 (P0407)